

**STable 2:** Antimicrobial susceptibility of streptococcal isolates cultured from the oropharynx of 88 individuals (children and their parents) from the general population of Belgium.

	Prevalence, n/N participants (%)	N isolates	Azithromycin, mg/L median (IQR)	P- value
<b>All Streptococcal spp.</b>	<b>88/88 (100)</b>	<b>312</b>	<b>1 (0.5-12)</b>	<b>0.6098</b>
Parents	38/38 (100)	128	<b>1 (0.5-12)</b>	
Children	50/50 (100)	181	<b>1 (0.5-12)</b>	
<b>S. mitis/oralis</b>	<b>82/88 (93.2)</b>	<b>146</b>	<b>1.25 (0.38-16)</b>	<b>0.5325</b>
Parents	35/38 (92.1)	68	1 (0.5-12)	
Children	47/50 (94.0)	78	4 (0.5-16)	
<b>S. salivarius</b>	<b>47/88 (53.4)</b>	<b>56</b>	<b>0.875 (0.5-12)</b>	<b>0.1115</b>
Parents	15/38 (46.9)	18	1 (0.75-12)	
Children	32/50 (64.0)	38	0.75 (0.5-8)	
<b>S. parasanguinis</b>	<b>28/88 (31.8)</b>	<b>30</b>	<b>8 (1-16)</b>	<b>0.3136</b>
Parents	12/36 (33.3)	14	8 (1-16)	
Children	16/50 (32.0)	16	10 (1-32)	
<b>S. vestibularis</b>	<b>24/88 (27.3)</b>	<b>29</b>	<b>1 (0.5-12)</b>	<b>0.6729</b>
Parents	11/36 (30.6)	13	0.75 (0.5-12)	
Children	13/50 (26.0)	16	2.5 (0.63-14)	
<b>S. sanguinis</b>	<b>17/88 (19.3)</b>	<b>19</b>	<b>0.25 (0.13-6)</b>	<b>0.6110</b>
Parents	8/36 (22.2)	8	0.345 (0.11-9)	
Children	9/50 (18.0)	11	0.25 (0.13-0.38)	
<b>S. pneumoniae</b>	<b>14/88 (15.9)</b>	<b>14</b>	<b>8 (0.38-16)</b>	<b>0.3317</b>
Parents	12/36 (33.3)	4	10 (0.75-16)	
Children	16/50 (32.0)	10	0.375 (0.25-12)	

IQR = interquartile range