

**Supplementary Table 2: Association between sociodemographic characteristics and medication adherence.**

Variables	Adherent Frequency (%)	Non-adherent Frequency (%)	$\chi^2$	P value	Exp (B)	95% CI
<b>N=410</b>						
<b>Age Group, years (Mean±SD: 31.58±9.47; 19-61=42)</b>						
19–28	76 (40.9)	110 (59.1)	8.60	0.072	1.192	0.924–1.540
29–38	31 (25.0)	93 (75.3)	-	-	-	-
39–48	29 (38.2)	47 (61.8)	-	-	-	-
49–58	7 (33.3)	14 (66.7)	-	-	-	-
≥59	1 (33.3)	2 (66.7)	-	-	-	-
<b>Sex</b>						
Male	76 (33.0)	154 (67.0)	0.993	0.319	0.878	0.552–1399
Female	68 (37.8)	112 (62.2)	-	-	-	-
<b>Educational Status*</b>						
No education	11 (45.8)	23 (54.2)	1.079	0.782	0.996	0.782–1.269
Low education	33 (34.7)	62 (65.3)	-	-	-	-
High education	44 (39.0)	69 (61.0)	-	-	-	-
Qur'anic education	56 (33.3)	112 (66.7)	-	-	-	-
<b>Occupational Status</b>						
Class I	1 (100.0)	0 (0.0)	2.33	0.674	0.955	0.440–2.112
Class II	18 (34.6)	34 (65.4)	-	-	-	-
Class III	41 (35.7)	74 (64.3)	-	-	-	-
Class IV	17 (31.0)	38 (69.0)	-	-	-	-
Class V	67 (35.8)	120 (64.2)	-	-	-	-
<b>Marital Status</b>						
Single	70 (37.2)	118 (62.8)	8.867	0.031 <sup>†</sup>	3.13	1.02–9.63
Married	54 (59.5)	129 (70.5)	-	-	-	-
Divorced	10 (43.5)	13 (56.5)	-	-	-	-
Widowed	10 (62.5)	6 (37.5)	-	-	-	-

\*High education is having ≥12 years of western education, low education <12 years.

<sup>†</sup>p<0.05

SD: standard deviation.